

金沢医療センター小児科業績集 (2020 年度)

※ 原著 (英文)

1) **Clinical Characteristics of Pediatric Pyelonephritis Without Pyuria or Bacteriuria.**

Yokoyama T, Takemura Y, Irabu H, Taniguchi C, Jinkawa A, Yamamiya M, Obata M, Miyashita K, Inoue M, Sakazume S, Ohta K.

Pediatr Infect Dis J. 2020. 39 (5); 385-388.

doi:10.1097/INF.0000000000002609.

2) **Apoptosis inhibitor of macrophage as a biomarker for disease activity in Japanese children with IgA nephropathy and Henoch-Schönlein purpura nephritis.**

Irabu H, Shimizu M, Kaneko S, Inoue N, Mizuta M, Tasaki Y, Ohta K, Yachie A, Wada T.

Pediatr Res. 2021 Feb; 89(3): 667-672.

doi: 10.1038/s41390-020-0951-1. Epub 2020 May 14.

3) **Clinical Significance of Serum Galactose-Deficient IgA1 Level in Children with IgA Nephropathy**

Irabu H, Shimizu M, Kaneko S, Inoue N, Mizuta M, Ohta K, Yachie A.

Journal of Immunology Research. 2020 May 21;2020:4284379.

doi: 10.1155/2020/4284379. eCollection 2020.

4) **Recurrent painful ophthalmoplegic neuropathy**

Yokoyama T, Yamamiya M, Takakuwa M, Taniguchi C, Koizumi E, Jinkawa A, Obata M, Miyashita K, Inoue M, Sakazume S, Ohta K
J Paediatr Child Health. 2020 Sep 1. doi: 10.1111/jpc.15147. Online ahead of print.

5) **Serum insulin-like growth factor-binding protein 2 levels as an indicator for disease severity in enterohemorrhagic *Escherichia coli* induced hemolytic uremic syndrome**

Tasaki Y, Inouea N, Shimizu M , Sugimoto N, Ishikawa I, Mizuta M, Yokoyama T, Kuroda M, Ohta K, Yachie A, Wada T

Renal Failure. 2021. 43 (1) ; 382–387

doi: 10.1080/0886022X.2021.1885445.